arXiv:1605.08640v1 [math.CO] 27 May 2016

ARITHMETICAL SEMIRINGS

MARCO ALDI

Abstract. In the context of abstract analytic number theory, we introduce the notion of arithmetical semiring. We show how the language of arithmetical semirings allows us to relate the asymptotic distributions additive and multiplicative primes. As an application we study the asymptotic distribution of connected cartesian-prime graphs among all graphs.

1. introduction

Let $\mathbb{G}$ be the set of all unlabeled finite simple graphs (here simply referred to as *graphs*), let $\mathbb{G}^+ \subseteq \mathbb{G}$ be the subset of all connected graphs and let $\mathbb{G}^\square \subseteq \mathbb{G}^+$ be the set of all *cartesian-prime graphs* i.e. the set of all connected graphs that cannot be written as the cartesian product of two graphs with fewer vertices. Moreover, let $\mathbb{G}(n)$ be the number of graphs with n vertices, let $\mathbb{G}^+(n)$ be the number of connected graphs with n vertices and let $\mathbb{G}^\square(n)$ be the number of cartesian-prime graphs with n vertices. As proved by Wright [11], $\mathbb{G}(n) \sim \mathbb{G}^+(n)$ i.e. almost all graphs are connected. In this paper we use this fact to give a short proof that

$$\mathbb{G}^+(n) - \mathbb{G}^\square(n) = O\left(\frac{2^{\binom{\lfloor n/2 \rfloor}{2}}}{\lfloor n/2 \rfloor!}\right). \tag{1}$$

In particular, (1) implies that $\mathbb{G}^+(n) \sim \mathbb{G}^\square(n)$ i.e. almost all connected graphs are cartesian-prime (a statement which appears to be well-known to experts, but that we were unable to locate in the literature). Furthermore, for graphs with an even number of vertices we obtain a more precise asymptotic expansion

$$\mathbb{G}^+(2n) - \mathbb{G}^\square(2n) = \frac{2^{\binom{n}{2}}}{n!} + \sum_{s=1}^{R-1} \omega_s(n) \frac{2^{\binom{n-s}{2}}}{(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right) \tag{2}$$

where R is an arbitrary integer greater than 1 and, for each integer $s \geq 1$, ω_s is a polynomial of degree s.

Enumerative formulas for the number of cartesian prime graphs were derived in [5] in the context of the theory of species. In this paper we derive asymptotic information in the context of abstract analytic number theory [4]. More precisely, we introduce the notion of *arithmetical semiring* which consists of an additive arithmetical semigroup endowed with a second compatible operation with respect to which the set of additive primes is a (multiplicative) arithmetical semigroup. While the notion of arithmetical semiring seems rather natural from the point of view of abstract analytic number theory, we were unable to find previous references to it in the literature. The connection between graph theory and abstract

analytic number theory was first observed in [3] where it was shown that the results of [10] and [11] can be used to study the statistics of arithmetical functions on arbitrary additive arithmetical semigroups with the property that almost all elements are (additive) primes.

In this paper we point out that the results of [3] can be further extended to arithmetical semirings. More precisely, our main theorem states that if an arithmetical semiring is *monotonic* (i.e. the number $\mathbb{S}^+(n)$ of additive primes of degree n is a non-decreasing function of n) then the number of degree n elements of $\mathbb{S}$ that are not multiplicative primes is $O\left(\mathbb{S}^+\left(\left\lfloor \frac{n}{2} \right\rfloor\right)\right)$. As a consequence, if almost all elements of $\mathbb{S}$ are additive primes (a condition named axiom $\mathcal{G}_1^{\#}$ in [3]) then almost all elements of $\mathbb{S}$ are multiplicative primes as well. As an application, we establishes analogues of (1) and (2) for the semiring $\mathbb{G}_{\text{ev}}$ of graphs with an even number of edges.

We conclude the paper with a study of submultiplicative functions on arithmetical semirings. In particular we prove that if f is a submultiplicative function on an monotonic arithmetical semiring in which almost all elements are additive primes, then the average of f on additive primes is asymptotic to the average of f on multiplicative primes. From this fact we derive multiplicative analogues of Knopfmacher's results regarding the asymptotic behavior of several arithmetical functions of importance in abstract analytic number theory.

Acknowledgments: The author would like to thank Richard Hammack for patiently answering his questions on the cartesian product of graphs.

2. Arithmetical Semirings

Definition 1. Let $(\mathbb{U}, \star)$ be a free commutative semigroup with identity $\mathbf{e}_\star$. We say that $(\mathbb{U}, \star)$ is a *unique factorization semigroup* if there exists a finite or countable free generating subset $\mathbb{U}^\star \subseteq U \setminus \{\mathbf{e}_\star\}$ whose elements are called $\star$-*primes* such that every $G \in U \setminus \{\mathbf{e}_\star\}$ can be uniquely (up to permutation of the factors) written as

$$G = G_{i_1} \star G_{i_2} \star \ldots \star G_{i_n}$$

for some (not necessarily distinct) $G_{i_1}, \ldots, G_{i_n} \in \mathbb{U}^\star$.

Definition 2. Given a unique factorization semigroup $(\mathbb{A}, +)$ and a semigroup homomorphism $\partial : (\mathbb{A}, +) \to (\mathbb{R}_{\geq 0}, +)$, let $\underline{\mathbb{A}}(x) = \partial^{-1}(x)$ and let $\mathbb{A}(x) = |\underline{\mathbb{A}}(x)|$ for each $x \in \mathbb{R}_{\geq 0}$. Similarly, we write $\underline{\mathbb{A}}^+(x) = \partial^{-1}(x) \cap \mathbb{A}^+$ and $\mathbb{A}^+(x) = |\underline{\mathbb{A}}^+(x)|$ for each $x \in \mathbb{R}_{\geq 0}$. We say that $(\mathbb{A}, +, \partial)$ is an *additive arithmetical semigroup* if

i) $\underline{\mathbb{A}}(0) = \{\mathbf{e}_+\}$;
ii) $\mathbb{A}(x)$ is finite for each $x \in \mathbb{R}_{>0}$;
iii) $\mathbb{A}(x) = 0$ unless x is an integer.

Example 3. Let $\mathbb{G}$ be the collection of all (unlabeled finite simple) graphs thought of as a semigroup with respect to the operation $+$ of disjoint union. If ∂ is the function that to each $G \in \mathbb{G}$ assigns the corresponding number of vertices $\partial(G) = |V(G)|$, then $(\mathbb{G}, +, \partial)$ is an additive arithmetical semigroup [3]. The set of additive primes $\mathbb{G}^+$ coincides by definition with the set of *connected graphs*.

Definition 4. Let $(\mathbb{S}, +, \square)$ be a commutative unital semiring and let $\partial : (\mathbb{S}, +, \square) \to (\mathbb{R}_{\geq 0}, +, \cdot)$ be a semiring homomorphism. We say that $(\mathbb{S}, +, \square, \partial)$ is an *arithmetical semiring* if $(\mathbb{S}, +, \partial)$ is an additive arithmetical semiring and $(\mathbb{S}^+, \square, \partial)$ is a unique factorization semigroup such that $\underline{\mathbb{S}}^+(1) = \{\mathbf{e}_\square\}$.

Remark 5. If $(\mathbb{S}, +, \square, \partial)$ is an arithmetical semiring, then in particular $(\mathbb{S}^+, \square, \partial)$ is an arithmetical semigroup in the sense of [4].

Remark 6. By unique factorization, the assignment $S \mapsto S + \mathbf{e}_\square$ defines an injection $\underline{\mathbb{S}}(n) \to \underline{\mathbb{S}}(n+1)$ for all n. In particular, the sequence $\{\mathbb{S}(n)\}$ is non-decreasing.

Definition 7. An arithmetical semiring $(\mathbb{S}, +, \square, \partial)$ is *monotonic* (resp. *strictly monotonic*) if there exists $N \in \mathbb{N}$ such that $\mathbb{S}^+(n) \leq \mathbb{S}^+(n+1)$ (resp. $\mathbb{S}^+(n) < \mathbb{S}^+(n+1)$).

Example 8. Let $(\mathbb{G}, +, \partial)$ be the additive arithmetical semigroup of graphs as in Example 3. The cartesian product of two graphs $G_1, G_2 \in \mathbb{G}$ is defined as the graph $G_1 \square G_2 \in \mathbb{G}$ with vertices $V(G_1 \square G_2) = V(G_1) \times V(G_2)$ and edges

$$E(G_1 \square G_2) = (E(G_1) \times \Delta_2) \cup (\Delta_1 \times E(G_2)) \,. \tag{3}$$

Here $\Delta_i \subseteq (V(G_i))^2$ is the diagonal and the right hand side of (3) is thought of as a subset of $(V(G_1 \square G_2))^2$ by means of the canonical identification of the latter with $(V(G_1))^2 \times (V(G_2))^2$. According to a theorem of Sabidussi and Vizing ([8], [9]), $(\mathbb{G}^+, \square)$ is a unique factorization semigroup. Moreover it is easy to check (see e.g. [2]) that $(\mathbb{G}, +, \square)$ is a semiring and ∂ is a semiring homomorphism. Since the only graph of degree 1 is the graph K_1 with only one vertex (the multiplicative identity with respect to $\square$), we conclude that $(\mathbb{G}, +, \square, \partial)$ is an arithmetical semiring. Since for each $G \in \underline{\mathbb{G}}^+(n)$, the only vertex of degree 0 in $G + K_1$ can be connected in at least one way to the vertices of G, then $(\mathbb{G}, +, \square, \partial)$ is monotonic.

Example 9. If $(\mathbb{S}, +, \square, \partial)$ is an arithmetical semiring and $\mathbb{S}' \subseteq \mathbb{S}$ is a subsemiring that contains $\mathbf{e}_+$ and $\mathbf{e}_\square$, then $(\mathbb{S}', +, \square, \partial)$ is an arithmetical semiring. For instance it follows from (3) that the subset $\mathbb{G}_{\text{ev}} \subseteq \mathbb{G}$ of graphs with an even number of edges is closed under the semiring operations defined in Example 8. Therefore, $(\mathbb{G}_{\text{ev}}, +, \square, \partial)$ is an arithmetical semiring. Another interesting subsemiring of $\mathbb{G}$ is the *Hamming semiring* $\mathbb{H} = \mathbb{N}[K_2, K_3, \ldots]$ generated by all complete graphs (with constants counting connected components isomorphic to K_1). Since for every prime number p one has $\underline{\mathbb{H}}(p) = \{K_p\}$ and $\underline{\mathbb{H}}(2p) = \{K_{2p}, K_2 \square K_p\}$, we conclude that the arithmetical semiring $(\mathbb{H}, +, \square, \partial)$ is not monotonic.

Example 10. Upon canonically identifying the Hamming semiring $\mathbb{H}$ with a semiring of polynomials in infinitely many variables over $\mathbb{N}$, the *Hamming polynomial* defined in [1] can be thought of as the semiring homomorphism $c : (\mathbb{G}, +\square) \to (\mathbb{H}, +, \square)$ that to each $G \in \mathbb{G}$ assigns

$$c(G) = \sum \alpha_{i_1, \ldots, i_m} K_{i_1} \square \cdots \square K_{i_m}$$

where $\alpha_{i_1, \ldots, i_m}$ (where the indices $i_1, \ldots, i_m$ are possibly repeated) is the number of induced subgraphs of G that are isomorphic to $K_{i_1} \square \cdots \square K_{i_m}$. Since $\partial(G) \leq \partial(c(G))$, we conclude that $(\mathbb{G}, +, \square, \partial \circ c)$ defines a non-standard structure of arithmetical semiring structure with

underlying semiring $(\mathbb{G}, +, \Box)$. More generally, one can consider arithmetical semirings of the form $(\mathbb{G}, +, \Box, f \circ c)$, for some semiring homomorphism $f : (\mathbb{H}, +, \Box) \to (\mathbb{N}, +, \cdot)$.

3. Knopfmacher Axioms

Definition 11. An arithmetical semiring $(\mathbb{S}, +, \Box, \partial)$ may additionally satisfy one or more axioms from the following list:

$\mathcal{G}_1^+$**:** $\mathbb{S}(n) \sim \mathbb{S}^+(n)$;
$\mathcal{G}_1^\Box$**:** $\mathbb{S}^+(n) \sim \mathbb{S}^\Box(n)$;
$\mathcal{G}_2^+$**:** $\mathbb{S}^+(n-1) = o(\mathbb{S}^+(n))$;
$\mathcal{G}_2^\Box$**:** $\mathbb{S}^\Box\left(\left\lfloor \frac{n}{2} \right\rfloor\right) = o(\mathbb{S}^\Box(n))$;
$\mathcal{G}_3^+$**:** $\mathcal{G}_2^+$ holds and $\mathbb{S}(n) - \mathbb{S}^+(n) = O(\mathbb{S}^+(n-1))$;
$\mathcal{G}_3^\Box$**:** $\mathcal{G}_2^\Box$ holds and $\mathbb{S}^+(n) - \mathbb{S}^\Box(n) = O\left(\mathbb{S}^\Box\left(\left\lfloor \frac{n}{2} \right\rfloor\right)\right)$.

Remark 12. Axioms $\mathcal{G}_i^+$, $k = 1, 2, 3$ depend only on the underlying structure of additive arithmetical semigroup and are equivalent to axioms $\mathcal{G}_i^\#$ defined in [3]. Axioms $\mathcal{G}_i^\Box$ are introduced here as their natural multiplicative analogues.

Remark 13. An arithmetical semiring $(\mathbb{S}, +, \Box, \partial)$ for which axiom $\mathcal{G}_2^+$ holds is in particular strictly monotonic. On the other hand, if $(\mathbb{S}, +, \Box, \partial)$ is strictly monotonic, then $n = O(\mathbb{S}^+(n))$.

Lemma 14. *Let $(\mathbb{S}, +, \Box, \partial)$ be an arithmetical semiring and suppose that there exist $\alpha, \beta, \gamma, a, b \in \mathbb{R}$ such that $\alpha > 0$, $a > b > 0$ and*

$$\log \mathbb{S}(n) = \alpha n^{a+1} + \beta n \log n + \gamma n + O(n^b)\,. \tag{4}$$

Then axioms $\mathcal{G}_1^+$ and $\mathcal{G}_2^+$ are satisfied. Moreover, if B is the integer sequence defined recursively for all positive integers by the formula

$$B(n) = -\mathbb{S}(n) - \sum_{s=1}^{n-1} B(s)\mathbb{S}(n-s)\,, \tag{5}$$

then

$$\mathbb{S}^+(n) = \mathbb{S}(n) + \sum_{s=1}^{R-1} B(s)\mathbb{S}(n-s) + O(\mathbb{S}(n-R)) \tag{6}$$

for each integer $R > 1$.

Proof: According to [3], the sequences $\{\mathbb{S}(n)\}$ and $\{\mathbb{S}^+(n)\}$ are related by the additive Euler product formula

$$\sum_{n=0}^{\infty} \mathbb{S}(n)x^n = \prod_{m=1}^{\infty} (1 - x^m)^{-\mathbb{S}^+(m)}$$

understood as an equality of formal power series. Since by assumption $\mathbb{S}(n) \geq \mathbb{S}^+(n) > 0$ for n sufficiently large, the result follows by Theorem 5 and Theorem 7 in [13]. □

Example 15. Let $(\mathbb{G}, +, \Box, \partial)$ be the arithmetical semiring of graphs introduced in Example 8. As shown in [12], the sequence $\{\mathbb{G}(n)\}$ has asymptotic expansion

$$\mathbb{G}(n) = \frac{2^{\binom{n}{2}}}{n!} + \sum_{s=1}^{R-1} \varphi_s(n) \frac{2^{\binom{n-s}{2}}}{(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right) \tag{7}$$

where R is any integer greater than 1 and, for each $s \in \mathbb{N}$, $\varphi_s(n)$ is polynomial of degree s in n. An explicit group-theoretic formula for φ_s is given in [12]. For small values of s it yields

$$\begin{aligned}
\varphi_1(n) &= n - 1\,; \\
\varphi_2(n) &= \frac{1}{3}(3n^2 - 13n + 14)\,; \\
\varphi_3(n) &= \frac{1}{3}(4n^3 - 46n^2 + 177n - 225)\,; \\
\varphi_4(n) &= \frac{2}{45}(60n^4 - 1305n^3 + 10580n^2 - 37737n + 49828)\,.
\end{aligned}$$

Therefore (4) holds with $\alpha = \log(\sqrt{2})$, $\beta = -1$, $\gamma = -\alpha - \beta$, $a = 1$ and, say, $b = \frac{1}{2}$. By Lemma 14, the arithmetical semiring $(\mathbb{G}, +, \Box, \partial)$ satisfies axioms $\mathcal{G}_1^+$ and $\mathcal{G}_2^+$. Substituting (7) into (6) one obtains the asymptotic expansion

$$\mathbb{G}^+(n) = \frac{2^{\binom{n}{2}}}{n!} + \sum_{s=1}^{R-1} \omega_s(n) \frac{2^{\binom{n-s}{2}}}{(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right) \tag{8}$$

found in [13], where R is any integer greater than 1 and for all $s \in \mathbb{N}$

$$\omega_s(n) = \varphi_s(n) + B(s) + \sum_{r=1}^{s-1} B(r)\varphi_s(n-r)$$

where B is defined by specializing (5) to $\mathbb{S} = \mathbb{G}$. For instance, using $\mathbb{G}(1) = 1$, $\mathbb{G}(2) = 2$, $\mathbb{G}(3) = 4$ and $\mathbb{G}(4) = 11$ one obtains $B(1) = B(2) = B(3) = -1$, $B(4) = -4$ and thus [13]

$$\begin{aligned}
\omega_1(n) &= n - 2\,; \\
\omega_2(n) &= \frac{1}{3}(3n^2 - 16n + 17)\,; \\
\omega_3(n) &= \frac{1}{3}(4n^3 - 49n^2 + 193n - 249)\,; \\
\omega_4(n) &= \frac{1}{45}(120n^4 - 2670n^3 + 21985n^2 - 79359n + 105656)\,.
\end{aligned}$$

Example 16. As observed in [6], $2\mathbb{G}_{\mathrm{ev}}(n) - \mathbb{G}(n)$ equals the number of self-complementary graphs with n vertices. Therefore, using the asymptotic estimates of [7], we obtain

$$\log(2\mathbb{G}_{\mathrm{ev}}(n) - \mathbb{G}(n)) = o\left(\log\left(\mathbb{G}\left(3\left\lfloor\frac{n}{2}\right\rfloor\right)\right)\right)$$

and thus

$$2\mathbb{G}_{\rm ev}(n) - \mathbb{G}(n) = O\left(\mathbb{G}\left(3\left\lfloor\frac{n}{2}\right\rfloor\right)\right) = O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right)$$

for all integers $R > 1$. Therefore

$$\mathbb{G}_{\rm ev}(n) = \frac{2^{\binom{n}{2}}}{2n!} + \sum_{s=1}^{R-1}\varphi_s(n)\frac{2^{\binom{n-s}{2}}}{2(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right)$$

where the polynomials φ_s are as in (7) and R is any integer greater than 1. In particular, the arithmetical semiring $(\mathbb{G}_{\rm ev}, +, \Box, \partial)$ satisfies the assumptions of Lemma 14. Thus, for any integer $R > 1$,

$$\mathbb{G}^+_{\rm ev}(n) = \frac{2^{\binom{n}{2}}}{2n!} + \sum_{s=1}^{R-1}\omega^{\rm ev}_s(n)\frac{2^{\binom{n-s}{2}}}{2(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right), \tag{9}$$

where

$$\omega^{\rm ev}_s(n) = \varphi_s(n) + B_{\rm ev}(s) + \sum_{r=1}^{s-1} B_{\rm ev}(r)\varphi_s(n-r)$$

for each $s \in \mathbb{N}$ and $B_{\rm ev}$ is the sequence obtained by specializing (5) to $\mathbb{S} = \mathbb{G}_{\rm ev}$. For instance, using $\mathbb{G}_{\rm ev}(1) = 1$, $\mathbb{G}_{\rm ev}(2) = 1$, $\mathbb{G}_{\rm ev}(3) = 2$, $\mathbb{G}_{\rm ev}(4) = 6$ we obtain $B(1) = 1$, $B(2) = 0$, $B(3) = -1$ and $B(4) = -3$ and thus

$$\begin{aligned}
\omega^{\rm ev}_1(n) &= n - 2\,;\\
\omega^{\rm ev}_2(n) &= \frac{1}{3}(3n^2 - 16n + 20)\,;\\
\omega^{\rm ev}_3(n) &= \frac{1}{3}(4n^3 - 49n^2 + 196n - 258)\,;\\
\omega^{\rm ev}_4(n) &= \frac{1}{45}(120n^4 - 2670n^3 + 22030n^2 - 79734n + 106481)\,.
\end{aligned}$$

Lemma 17. *Let $(\mathbb{S}, +, \Box, \partial)$ be an arithmetical semiring. Then for each $n \in \mathbb{N}$,*

$$\sum_{s=1}^{n-1}\mathbb{S}^+(s)\mathbb{S}^+(n-s) \le 2(\mathbb{S}(n) - \mathbb{S}^+(n)) - \left(\mathbb{S}^+\left(\frac{n}{2}\right)\right)^2 \le \sum_{s=1}^{n-1}\mathbb{S}(s)\mathbb{S}(n-s) \tag{10}$$

and

$$\sum_{r=2}^{\lfloor n/2\rfloor}\mathbb{S}^\Box(r)\mathbb{S}^\Box(n/r) \le 2(\mathbb{S}^+(n) - \mathbb{S}^\Box(n)) - \left(\mathbb{S}^\Box(\sqrt{n})\right)^2 \le \sum_{r=2}^{\lfloor n/2\rfloor}\mathbb{S}^+(r)\mathbb{S}^+(n/r)\,. \tag{11}$$

Proof: $\mathbb{S}(n) - \mathbb{S}^+(n)$ is the number of elements of $\underline{\mathbb{S}}(n)$ that are not $+$-primes. On the one hand, this is greater or equal than the number of elements of $\underline{\mathbb{S}}(n)$ that can be written as

sum of exactly two +-primes i.e.

$$\frac{1}{2}\sum_{s=1}^{n-1}\mathbb{S}^+(s)\mathbb{S}^+(n-s)+\frac{1}{2}\left(\mathbb{S}^+\left(\frac{n}{2}\right)\right)^2\leq \mathbb{S}(n)-\mathbb{S}^+(n)\,.$$

On the other hand, each element of $\underline{\mathbb{S}}(n)$ that is not a prime is also in $\underline{\mathbb{S}}(s)+\underline{\mathbb{S}}(n-s)$ for some (possibly several) $s\in\{1,\dots,n-1\}$. Therefore,

$$\mathbb{S}(n)-\mathbb{S}^+(n)\leq \frac{1}{2}\sum_{s=1}^{n-1}\mathbb{S}(s)\mathbb{S}(n-s)+\frac{1}{2}\left(\mathbb{S}^+\left(\frac{n}{2}\right)\right)^2\,.$$

The last term corrects the possible undercounting of elements of $2\underline{\mathbb{S}}(n/2)$ since elements of $2\underline{\mathbb{S}}(n/2)\setminus\mathbb{S}^+$ also belong to $\underline{\mathbb{S}}(s)+\underline{\mathbb{S}}(n-s)$ for some $s<n/2$. A similar argument proves (11). □

Lemma 18. *Let $(\mathbb{S},+,\square,\partial)$ be a monotonic arithmetical semiring and let n,D be positive integers such that $2<D<\sqrt{n}$. Then*

$$\begin{aligned}\mathbb{S}^+(n)-\mathbb{S}^\square(n) &\leq \frac{1}{2}\left(\mathbb{S}^\square(\sqrt{n})\right)^2+\sum_{r=2}^{D-1}\mathbb{S}^+(r)\mathbb{S}^+\left(\frac{n}{r}\right)+\\ &+\frac{1}{2}\sum_{r=D}^{\lfloor n/D\rfloor}\mathbb{S}^+(r)\mathbb{S}^+\left(\left\lfloor\frac{n}{D}\right\rfloor+D-r\right)\,.\end{aligned}$$

Proof: Let r be a divisor of n such that $D\leq r\leq\lfloor n/D\rfloor$. Then $n\geq rD$ and thus $n\left(\frac{1}{D}-\frac{1}{r}\right)\geq(r-D)$ which implies

$$\frac{n}{r}\leq\left\lfloor\frac{n}{D}\right\rfloor+D-r\,. \tag{12}$$

It follows from (11) that

$$\mathbb{S}^+(n)-\mathbb{S}^\square(n)\leq\frac{1}{2}\left(\mathbb{S}^\square(\sqrt{n})\right)^2+\sum_{r=2}^{D-1}\mathbb{S}^\square(r)\mathbb{S}^\square(n/r)+\frac{1}{2}\sum_{r=D}^{\lfloor n/D\rfloor}\mathbb{S}^\square(r)\mathbb{S}^\square(n/r)\,. \tag{13}$$

On the other hand

$$\mathbb{S}^\square(r)\mathbb{S}^\square(n/r)\leq\mathbb{S}^+(r)\mathbb{S}^+(n/r)\leq\mathbb{S}^+(r)\mathbb{S}^+\left(\left\lfloor\frac{n}{D}\right\rfloor+D-r\right) \tag{14}$$

where (12) and monotonicity were using to obtain the second inequality. The result then follows by substituting (14) into (11). □

Theorem 19. *If $(\mathbb{S},+,\square,\partial)$ is a monotonic arithmetical semiring then*
1) $\mathbb{S}^+(n)-\mathbb{S}^\square(n)=O\left(\mathbb{S}\left\lfloor\frac{n}{2}\right\rfloor\right)$;
2) axiom $\mathcal{G}_1^+$ implies axiom $\mathcal{G}_1^\square$.

Proof: Since

$$\frac{1}{2}\left(\mathbb{S}^{\square}(\sqrt{n})\right)^2 \leq \frac{1}{2}\left(\mathbb{S}^{+}(\sqrt{n})\right)^2 \leq \mathbb{S}(2\sqrt{n})\,,$$

setting $D=3$ in Lemma 18 and using (10) we obtain

$$\mathbb{S}^{+}(n)-\mathbb{S}^{\square}(n) \leq \mathbb{S}(2\sqrt{n})+\mathbb{S}^{+}(2)\mathbb{S}^{+}\left(\left\lfloor\frac{n}{2}\right\rfloor\right)+\mathbb{S}\left(\left\lfloor\frac{n}{3}\right\rfloor+3\right)-\mathbb{S}^{+}\left(\left\lfloor\frac{n}{3}\right\rfloor+3\right)\,. \tag{15}$$

By Remark 6, each term on the right hand side of (15) is $O\left(\mathbb{S}\left\lfloor\frac{n}{2}\right\rfloor\right)$, which proves 1). If axiom $\mathcal{G}_1^{+}$ holds, (11) implies that the right hand side of (15) is $o\left(\mathbb{S}\left\lfloor\frac{n}{2}\right\rfloor+2\right)$. Using Remark 6 again, we obtain $\mathbb{S}^{+}(n)-\mathbb{S}^{\square}(n)=o(\mathbb{S}(n))$ and thus, by axiom $\mathcal{G}_1^{+}$, $\mathcal{G}_1^{\square}$ holds. □

Corollary 20. *If $(\mathbb{S},+,\square,\partial)$ is an arithmetical semiring that satisfies axioms $\mathcal{G}_1^{+}$ and $\mathcal{G}_2^{+}$ then it also satisfies axioms $\mathcal{G}_1^{\square}$, $\mathcal{G}_2^{\square}$ and $\mathcal{G}_3^{\square}$.*

Proof: By Remark 13 $(\mathbb{S},+,\square,\partial)$ is monotonic and thus, by Theorem 19, axiom $\mathcal{G}_1^{\square}$ holds. On the other hand, using (10) and $\mathcal{G}_1^{+}$ we obtain

$$\mathbb{S}^{\square}\left(\left\lfloor\frac{n}{2}\right\rfloor\right) \leq \mathbb{S}^{+}(2)\mathbb{S}^{+}\left(\left\lfloor\frac{n}{2}\right\rfloor\right) \leq \mathbb{S}\left(\left\lfloor\frac{n}{2}\right\rfloor+2\right)-\mathbb{S}^{+}\left(\left\lfloor\frac{n}{2}\right\rfloor+2\right)=o\left(\mathbb{S}\left(\left\lfloor\frac{n}{2}\right\rfloor+2\right)\right) \tag{16}$$

and thus, by monotonicity, $\mathcal{G}_2^{\square}$ holds. Since $\mathcal{G}_1^{+}$ and $\mathcal{G}_1^{\square}$ imply $\mathbb{S}(n)\sim\mathbb{S}^{\square}(n)$, axiom $\mathcal{G}_3^{\square}$ follows from Theorem 19. □

Corollary 21. *If $(\mathbb{S},+,\square,\partial)$ is a monotonic arithmetical semiring such that axiom $\mathcal{G}_1^{+}$ holds and $\mathbb{S}^{+}(2)=1$ then*

$$\mathbb{S}^{+}(2n)-\mathbb{S}^{\square}(2n)=\mathbb{S}^{+}(n)+o\left(\mathbb{S}^{+}\left(\left\lfloor\frac{2n}{3}\right\rfloor+3\right)\right)\,. \tag{17}$$

Proof: Substituting $\mathbb{S}^{\square}(2)=\mathbb{S}^{+}(2)=1$ in (11) we obtain $\mathbb{S}^{\square}(n)\leq\mathbb{S}^{+}(2n)-\mathbb{S}^{\square}(2n)$. Therefore, combining (15) and axiom $\mathcal{G}_1^{+}$ we obtain

$$\mathbb{S}^{+}(2n)-\mathbb{S}^{\square}(2n)-\mathbb{S}^{\square}(n)=\mathbb{S}^{+}(n)-\mathbb{S}^{\square}(n)+o\left(\mathbb{S}^{+}\left(\left\lfloor\frac{2n}{3}\right\rfloor+3\right)\right)\,.$$

The results follows by using axiom $\mathcal{G}_1^{\square}$, whose validity is guaranteed by Theorem 19. □

Remark 22. If $(\mathbb{S},+,\square,\partial)$ is an arithmetical semiring such that (4) holds and $\mathbb{S}^{+}(2)=1$, then combining (6) and (17) we obtain the asymptotic expansion

$$\mathbb{S}^{+}(2n)-\mathbb{S}^{\square}(2n)=\mathbb{S}(n)+\sum_{s=1}^{R-1}B(s)\mathbb{S}(n-s)+O(\mathbb{S}(n-R))\,, \tag{18}$$

where B is defined by (5).

Example 23. For the arithmetical semiring $(\mathbb{G},+,\square,\partial)$ of Example 8, Theorem 19 implies (1) and in particular $\mathcal{G}_1^{\square}$ holds i.e. almost all graphs are both connected and cartesian primes.

Furthermore, direct substitution of (8) into (17) yields (2). Similarly, Theorem 19 applied to the arithmetical semiring $(\mathbb{G}_{\rm ev}, +, \Box, \partial)$ of Example 9 yields

$$\mathbb{G}^{+}_{\rm ev}(n) - \mathbb{G}^{\Box}_{\rm ev}(n) = O\left(\frac{2^{\binom{\lfloor n/2 \rfloor}{2}}}{n!}\right).$$

Moreover, (17) implies that for each integer $R > 1$

$$\mathbb{G}^{+}_{\rm ev}(2n) - \mathbb{G}^{\Box}_{\rm ev}(2n) = \frac{2^{\binom{n}{2}}}{2n!} + \sum_{s=1}^{R-1} \omega^{\rm ev}_s(n) \frac{2^{\binom{n-s}{2}}}{2(n-s)!} + O\left(\frac{2^{\binom{n-R}{2}}}{(n-2R)!}\right)$$

where $\{\omega^{\rm ev}_s\}$ is the sequence of polynomials defined in Example 16.

4. submultiplicative functions

Definition 24. Let $(\mathbb{S}, +, \Box, \partial)$ be an arithmetical semiring. A function $f : \mathbb{S}^{+} \to \mathbb{R}_{\geq 0}$ is *submultiplicative* if $f(S_1 \Box S_2) \leq f(S_1) f(S_2)$ for all $S_1, S_2 \in \mathbb{S}^{+}$. Given f submultiplicative, and $\bullet \in \{+, \Box\}$ we define sequences $f^{\bullet}$, $f^{\bullet}_{\max}$ such that

$$f^{\bullet}(n) = \sum_{S \in \underline{\mathbb{S}}^{\bullet}(n)} f(S) \quad \text{and} \quad f^{\bullet}_{\max}(n) = \max_{s \leq n} \max_{S \in \underline{\mathbb{S}}^{\bullet}(n)} f(S)$$

for all $n \in \mathbb{N}$. We say that f has *asymptotic-mean value* μ on $\mathbb{S}^{\bullet}$ if

$$\lim_{n \to \infty} \frac{f^{\bullet}(n)}{\mathbb{S}^{\bullet}(n)} = \mu \,.$$

If this is the case, we say that f has asymptotic variance ν on $\mathbb{S}^{\bullet}$ if

$$\lim_{n \to \infty} \frac{1}{\mathbb{S}^{\bullet}(n)} \sum_{S \in \underline{\mathbb{S}}^{\bullet}(n)} (f(S) - \mu)^2 = \nu \,.$$

Theorem 25. *Let f be a submultiplicative function on the monotonic arithmetical semiring $(\mathbb{S}, +, \Box, \partial)$ such that $f^{+}_{\max}(n) = O(\mathbb{S}^{+}(n))$. Then*

$$f^{+}(n) - f^{\Box}(n) = O(\mathbb{S}(n) - \mathbb{S}^{+}(n)) \,. \tag{19}$$

Proof: Since the elements of $\mathbb{S}^{+}$ that are not $\Box$-primes contain at least one nontrivial $\Box$-prime factor, then

$$0 \leq f^{+}(n) - f^{\Box}(n) \leq \sum_{r=2}^{\lfloor n/2 \rfloor} f^{\Box}_{\max}(r) \mathbb{S}^{\Box}(r) f^{+}_{\max}\left(\frac{n}{r}\right) \mathbb{S}^{+}\left(\frac{n}{r}\right). \tag{20}$$

Separating the contribution from $r = 2$ and $r = 3$ on the right hand side of (20) and observing that $f^{\Box}_{\max}(n) \leq f^{+}_{\max}(n)$ for all $n \in \mathbb{N}$ we obtain

$$f^{+}(n) - f^{\Box}(n) = O\left(\mathbb{S}^{+}\left(\left\lfloor \frac{n}{2} \right\rfloor\right) \mathbb{S}^{+}\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + \sum_{r=4}^{\lfloor n/4 \rfloor} \left(\mathbb{S}^{+}(r) \mathbb{S}^{+}\left(\left\lfloor \frac{n}{4} \right\rfloor + 4 - r\right)\right)^2\right). \tag{21}$$

The term $\left(\mathbb{S}^+(r)\mathbb{S}^+\left(\left\lfloor\frac{n}{4}\right\rfloor+4-r\right)\right)^2$ can be thought of as counting elements of $\underline{\mathbb{S}}^+(n)$ whose $+$-prime factorization is the sum of two (not necessarily distinct) elements of $\underline{\mathbb{S}}^+(r)$, two (not necessarily distinct) elements of $\underline{\mathbb{S}}^+\left(\left\lfloor\frac{n}{4}\right\rfloor+4-r\right)$ and $n-2\left\lfloor\frac{n}{4}\right\rfloor-8$ copies of $\mathbf{e}_\square$. Therefore

$$\sum_{r=4}^{\lfloor n/4\rfloor}\left(\mathbb{S}^+(r)\mathbb{S}^+\left(\left\lfloor\frac{n}{4}\right\rfloor+4-r\right)\right)^2=O\left(\mathbb{S}(n)-\mathbb{S}^+(n)\right) \tag{22}$$

and thus (19) follows by substituting (22) into (21) and using (10). □

Corollary 26. *Let $(\mathbb{S},+,\square,\partial)$ be a monotonic arithmetical semiring that satisfies axiom $\mathcal{G}_1^+$ and let f be a submultiplicative function on $(\mathbb{S},+,\square,\partial)$ that has asymptotic mean-value μ on $\mathbb{S}^\square$. Then*

1) $f^+_{\max}(n)=O(\mathbb{S}^+(n))$ implies that f has asymptotic mean-value μ on $\mathbb{S}^+$;
2) $f^+_{\max}(n)=O\left(\mathbb{S}^+\left(\left\lfloor\frac{n}{2}\right\rfloor\right)\right)$ implies that f has asymptotic variance 0 on $\mathbb{S}^+$ if it has asymptotic variance 0 on $\mathbb{S}^\square$.

Proof: Axiom $\mathcal{G}_1^+$ applied to (19) implies $f^+(n)-f^\square(n)=o(\mathbb{S}^+(n))$. Therefore, using axiom $\mathcal{G}_1^+$ once more

$$\mu=\lim_{n\to\infty}\frac{f^\square(n)}{\mathbb{S}^\square(n)}=\lim_{n\to\infty}\frac{f^\square(n)}{\mathbb{S}^+(n)}=\lim_{n\to\infty}\frac{f^+(n)}{\mathbb{S}^+(n)}\,.$$

If $f^+_{\max}(n)=O\left(\mathbb{S}^+\left(\left\lfloor\frac{n}{2}\right\rfloor\right)\right)$, then $(f^2)^+_{\max}=O(\mathbb{S}^+(n))$. By assumption

$$0=\lim_{n\to\infty}\frac{1}{\mathbb{S}^\bullet(n)}\sum_{S\in\underline{\mathbb{S}}(n)}(f(S)-\mu)^2=\lim_{n\to\infty}\frac{(f^2)^\bullet(n)-2\mu f^\bullet(n)+\mu^2\mathbb{S}^\bullet(n)}{\mathbb{S}^\bullet(n)} \tag{23}$$

holds for $\bullet=\square$. Therefore, f^2 has asymptotic mean-value μ^2 on $\mathbb{S}^\square$ and thus, by Theorem 25, on $\mathbb{S}^+$. This implies that (23) holds for $\bullet=+$, which proves 2). □

Example 27. Let $(\mathbb{S},+,\square,\partial)$ be a monotonic arithmetical semiring for which $\mathcal{G}_1^+$ holds. The *divisor function* $d:\mathbb{S}^+\to\mathbb{R}_{\geq 0}$ that to each $S\in\mathbb{S}^+$ assigns the total number $d(S)$ of factorizations of the form $S=S'\square S''$ for some $S',S''\in\mathbb{S}^+$. Since $d(S)=2$ for all $S\in\mathbb{S}^\square$

$$\begin{aligned}
d^+_{\max}(n) &\leq 2+\sum_{r=2}^{\lfloor n/2\rfloor}\mathbb{S}^+(r)\\
&\leq 2+\mathbb{S}^+(2)+\mathbb{S}^+\left(\left\lfloor\frac{n}{2}\right\rfloor\right)+\sum_{r=3}^{\lfloor n/3\rfloor}\mathbb{S}^+(r)\mathbb{S}^+\left(\left\lfloor\frac{n}{3}\right\rfloor+3-r\right)\\
&= 2+\mathbb{S}^+(2)+O\left(\mathbb{S}^+\left(\left\lfloor\frac{n}{2}\right\rfloor\right)\right).
\end{aligned}$$

If we additionally assume that the sequence $\{\mathbb{S}^+(n)\}$ is unbounded (which is the case if axiom the arithmetical semiring is strictly monotone), then $d^+_{\max}(n)=O\left(\mathbb{S}^+\left(\left\lfloor\frac{n}{2}\right\rfloor\right)\right)$ and thus d satisfy the assumptions of Corollary 26. We conclude that d has asymptotic mean-value 2 and asymptotic variance 0 on $\mathbb{S}^+$.

Example 28. Let $(\mathbb{S}, +, \Box, \partial)$ be a strictly monotonic arithmetical semiring for which axiom $\mathcal{G}_1^+$ holds. Let $d_* : \mathbb{S}^+ \to \mathbb{R}_{\geq 0}$ be the *unitary-divisor function* that to each $S \in \mathbb{S}^+$ assigns the number $d_*(S)$ of factorizations $S = S' \Box S''$ for some $S', S'' \in \mathbb{S}^+$ that are $\Box$-*coprime* i.e. such that S' and S'' have no common $\Box$-prime factors. Since $d_*(S) \leq d(S)$ for each $S \in \mathbb{S}^+$ and thus $(d_*)^+_{\max} \leq d^+_{\max}$, then the argument of Example 27 applies to d_* as well. In particular d_* has asymptotic mean value 2 and asymptotic variance 0. Similarly, let $\beta : \mathbb{S}^+ \to \mathbb{N}$ be the *prime-divisor function* defined by the formula

$$\beta(S_1^{\Box\alpha_1} \Box S_2^{\Box\alpha_2} \Box \cdots \Box S_m^{\Box\alpha_m}) = \alpha_1\alpha_2 \cdots \alpha_m$$

for all $S_1, \ldots, S_m \in \mathbb{S}^\Box$ and non-negative integers $\alpha_1, \ldots, \alpha_m$. Then $\beta(S) \leq d(S)$ for each $S \in \mathbb{S}^+$, and thus β has asymptotic mean value 1 and asymptotic variance 0.

Example 29. Let $(\mathbb{S}, +, \Box, \partial)$ be a strictly monotonic arithmetical semiring for which $\mathcal{G}_1^+$ holds. Let $\sigma_* : \mathbb{S}^+ \to \mathbb{R}_{\geq 0}$ be the *divisor-sum function* defined by

$$\sigma_*(S) = \sum_{S = S' \Box S''} \partial(S')$$

for all $S \in \mathbb{S}^+$. Since

$$\sigma_*(S \Box S') = \sigma_*(S) + \sigma_*(S') \leq \sigma_*(S)\sigma_*(S')$$

for all $\Box$-coprime $S, S' \in \mathbb{S}^+$ and

$$\sigma_*(S^{\Box\alpha}) = 1 + \partial(S) + \cdots + (\partial(S))^\alpha \leq (1 + \partial(S))^\alpha = (\sigma_*(S))^\alpha$$

for all $S \in \mathbb{S}^\Box$, $\alpha \in \mathbb{N}$, then σ_* is submultiplicative. Moreover, Remark implies 13 $n = O\left((\mathbb{S}^+\left(\left\lfloor \frac{n}{2} \right\rfloor\right)\right)$. Using the function d introduced in Example 27, we arrive at

$$(\sigma_*)^+_{\max}(n) \leq (n+1)d^+_{\max}(n) = O(\mathbb{S}^+(n))\,.$$

Since σ_* has constant value $n + 1$ on $\underline{\mathbb{S}}^\Box(n)$, Theorem 25 implies

$$\sigma_*^+(n) = (n+1)\mathbb{S}^\Box(n) + o(\mathbb{S}^+(n))$$

and thus $\sigma_*^+(n) \sim (n+1)\mathbb{S}^+(n)$.

Example 30. Let $(\mathbb{S}, +, \Box, \partial)$ be a strictly monotonic arithmetical semiring for which axiom $\mathcal{G}_1^+$ holds. Let $\phi_* : \mathbb{S}^+ \to \mathbb{R}_{\geq 0}$ be the *Euler-type function* that to each $S \in \underline{\mathbb{S}}^+(n)$ assigns the number $\phi_*(S)$ of elements of $\underline{\mathbb{S}}^+(n)$ that are $\Box$-coprime to S. It is easy to see that ϕ_* is submultiplicative and that ϕ_* restricts to the constant function $\mathbb{S}^+(n) - 1$ on $\underline{\mathbb{S}}^\Box(n)$. Since $\mathbb{S}^+(n) - 1$ is also the maximum of ϕ_* on $\underline{\mathbb{S}}^+(n)$, we conclude from Theorem 25 that

$$\frac{\phi_*(n)}{\mathbb{S}^+(n)} \sim \mathbb{S}^+(n) - 1 \sim \mathbb{S}^+(n)\,.$$

References

[1] Boštjan Brešar, Paul Dorbec, Sandi Klavžar, and Michel Mollard, *Hamming polynomials and their partial derivatives*, European J. Combin. **28** (2007), no. 4, 1156–1162.

[2] Richard Hammack, Wilfried Imrich, and Sandi Klavžar, *Handbook of product graphs*, 2nd ed., Discrete Mathematics and its Applications (Boca Raton), CRC Press, Boca Raton, FL, 2011.

[3] J. Knopfmacher, *Arithmetical properties of finite graphs and polynomials*, J. Combinatorial Theory Ser. B **20** (1976), no. 3, 205–215.

[4] John Knopfmacher, *Abstract analytic number theory*, 2nd ed., Dover Books on Advanced Mathematics, Dover Publications, Inc., New York, 1990.

[5] Ji Li, *Prime graphs and exponential composition of species*, J. Combin. Theory Ser. A **115** (2008), no. 8, 1374–1401.

[6] Valery A. Liskovets, *Some easily derivable integer sequences*, J. Integer Seq. **3** (2000), no. 2, Article 00.2.2, 15.

[7] Edgar M. Palmer, *Asymptotic formulas for the number of self-complementary graphs and digraphs*, Mathematika **17** (1970), 85–90.

[8] Gert Sabidussi, *Graph multiplication*, Math. Z. **72** (1959/1960), 446–457.

[9] V. G. Vizing, *The cartesian product of graphs*, Vyčisl. Sistemy No. **9** (1963), 30–43 (Russian).

[10] E. M. Wright, *A relationship between two sequences*, Proc. London Math. Soc. (3) **17** (1967), 296–304.

[11] _____, *A relationship between two sequences. III*, J. London Math. Soc. **43** (1968), 720–724.

[12] _____, *The number of graphs on many unlabelled nodes*, Math. Ann. **183** (1969), 250–253.

[13] _____, *Asymptotic relations between enumerative functions in graph theory.*, Proc. London Math. Soc. (3) **20** (1970), 558–572.